

# N1-Arylation of 1,4-Benzodiazepine-2-ones with Diaryliodonium Salts


Raysa Khan<sup>a,1</sup>Robert Felix<sup>b</sup>Paul D. Kemmitt<sup>c</sup>Simon J. Coles<sup>d</sup>Graham J. Tizzard<sup>d</sup>John Spencer<sup>\*a</sup>

<sup>a</sup> Department of Chemistry, School of Life Sciences, University of Sussex, Falmer, BN19QJ, U.K.

<sup>b</sup> Tocris Bioscience, Tocris House, IO Centre, Moorend Farm Avenue, Bristol, BS11 0QL, U.K.

<sup>c</sup> Oncology, AstraZeneca, 310 Cambridge Science Park, Milton Road, Cambridge CB4 0WG, U.K.

<sup>d</sup> UK National Crystallography Service, School of Chemistry, University of Southampton, Highfield, Southampton, SO17 1BJ, U.K.

[j.spencer@sussex.ac.uk](mailto:j.spencer@sussex.ac.uk)

Received:

Accepted:

Published online:


DOI:

**Abstract** A library of *N*1-arylated 5-phenyl-1,3-dihydro-2*H*-1,4-benzodiazepin-2-ones has been synthesized starting with unsymmetrical diaryliodonium salts using aqueous ammonia as a base. This can also be applied to a similar 1,3,4-benzotriazepin-2-one derivative.

**Key words** Benzodiazepines, *N*-arylation, iodonium salt, privileged scaffold, benzotriazepine.

Compounds containing a 1,4-benzodiazepine scaffold are often termed as 'privileged structures' and are of significant interest to organic and medicinal chemists.<sup>1-18</sup> Many bioactive 1,4-benzodiazepines include *N*-arylated benzodiazepines; for example, the benzodiazepine derivative, **A** (Figure 1), is a bradykinin antagonist<sup>19</sup> and the related benzotriazepine **B** is an antagonist at the parathyroid hormone (PTH)-1 receptor.<sup>20</sup> Typically *N*-arylated benzodiazepines can be prepared by transition-metal catalysed couplings, often with copper, with various arylating agents. Generally, the reaction scope is limited with these routes and often require high temperatures and strong bases.<sup>19,21-23</sup>

Being able to generate libraries of diverse analogues, in this case by adding *N*-functionality to a privileged core unit, using mild and efficient methodologies, can substantially improve SAR studies (structure – activity relationship) and optimise the drug development process potentially repurposing privileged scaffolds for new biological targets.<sup>24,25</sup>



**Figure 1** Bioactive *N*-arylated benzodiazepine and benzotriazepine

We have an active interest in benzodiazepines<sup>26,27</sup> and recently reported a method to functionalise 5-phenyl-1,3-dihydro-2*H*-1,4-benzodiazepin-2-ones via a late-stage palladacycle assisted ortho C-H activation protocol.<sup>28,29</sup> Herein we present our approach to generate a series of *N*1-arylated 1,4-benzodiazepines using diaryliodonium salts. The latter react with nucleophiles in the absence of transition metal catalysts and are commonly used in organic synthesis as electrophilic reagents.<sup>30-35</sup>

Novak et al. recently reported a protocol for the *N*-arylation of pyrazoles.<sup>36</sup> A quick screen of conditions, adapting this protocol using diaryliodonium salts with weak bases under mild conditions, showed that it was indeed possible to perform similar arylations on the 1,4-benzodiazepine system. Upon initial screening of a number of solvents, DCE (1,2-dichloroethane) was found to give the best results (Table 1, entry 2). Solvents such as PEG (polypropylene glycol) and acetic acid gave poor yields. Similar results were observed on pyrazoles by Novak et al. where apolar solvents, immiscible in water, produced the best results.

**Table 1** Optimization of *N*-arylation of 1,4-benzodiazepines – solvent effects

Entry	Solvent	Conversion % <sup>a</sup>
1	Toluene	95
2	DCE	99
3	PEG	-
4	AcOH	-
5	Chloroform	85

<sup>a</sup> LCMS conversion.


A number of bases were tested subsequently and both NH<sub>3</sub> (25% w/w) and NaOH (sat. aq.) gave similar and the best results (Table 2, entry 1, 2).

**Table 2** Optimization of *N*-arylation of 1,4-benzodiazepines – base effects

Entry	Base	Conversion % <sup>a</sup>
1	NaOH (sat. aq.)	99
2	NH <sub>3</sub> (25% w/w)	99
3	K <sub>2</sub> CO <sub>3</sub>	80
4	NaH	-

<sup>a</sup> LCMS conversion.

Hence, optimal condition appeared to use NH<sub>3</sub> (aq.), DCE at room temperature for 30 min. Next, a series of functionalized 1,4-benzodiazepines was *N*-arylated using (4-nitrophenyl)phenyliodonium triflate in good to excellent yields (Scheme 1). Generally, in transition-metal-free processes unsymmetrical diaryliodonium salts give a mixture of products where both groups are transferred and the transfer of more sterically hindered and electron withdrawing group is preferable.<sup>34</sup> However, in this case (Scheme 1) only the nitrophenyl-group was transferred. We were able to *N*-arylate quite sterically hindered benzodiazepines such as **3e**, **3f** and **3g**. Of note, **3e** is a key intermediate towards **A**, the bradykinin B<sub>2</sub> receptor antagonist. We were also pleased to be able to conduct *N*-arylation on a previously ortho-arylated hindered benzodiazepine, **3h**, in good yield, whose structure was also confirmed by X-ray crystallography. Such molecules may be useful precursors to e.g., alpha-helical mimetics in medicinal chemistry.<sup>37,38</sup>


**Scheme 1** *N*-arylated 1,4-benzodiazepines


The use of other unsymmetrical diaryliodonium triflates was also explored (Table 3), which required longer reaction time and led to both aryl groups being transferred to obtain **3i** – **3l**. As expected, the transfer of more sterically hindered or less electron rich groups was preferred. Further attempts to use unsymmetrical diaryliodonium salts such as phenyl(3-methylphenyl)iodonium triflate, phenyl(4-methylphenyl)iodonium triflate and (2-methylphenyl)(2,4,6-trimethylphenyl)iodonium triflate gave little or no products. Additionally, attempted *N*-arylation with symmetrical diaryliodonium triflates or tetrafluoroborates such as bis(2-fluorophenyl)iodonium tetrafluoroborate and bis(4-bromophenyl)iodonium triflate, gave, at best, traces of products.

**Table 3** Further *N*-arylation of 1,4-benzodiazepines

Salt	Product (major)	Product (minor)
 4	 3i; 51%	 3j; 8%
 5	 3k; 42%	 3l; 9%

Reaction time = 8 h.

We have briefly explored the *N*-arylation on a 1,3,4-benzotriazepine, **6**, which resulted in di-arylation and yielded **7** (Scheme 2).



Scheme 2 N-arylation on a 1,3,4-benzotriazepine

Interestingly, the iodonium salts were observed to undergo reaction with water present in the reaction to give diarylether products. The ether product is only observed in substantial amounts when the benzodiazepine substrates react poorly with the diaryliodonium salts (Table 4). The ether product **10** was also obtained merely by stirring the iodonium salt with water in DCE with a mild base for 20 min at room temperature with a yield of 43%. Olofsson et al. have reported the synthesis of related diarylethers by reacting diaryliodonium salts with phenols in the presence of mild bases.<sup>39</sup>

Table 4 Diaryl ether formation

Substrate	Expected product	Observed product
 8	 9; 0%	 10; 43%
 11	 12; 0%	 10; 45%

In summary we have presented a mild metal-free route to N-arylated benzodiazepines, three of which were structurally characterized in the solid state (**3a**, **3h**, **3i**, with CCDC numbers 1560492-1560494 respectively.<sup>40</sup>

## Experimental Section

All reactions were conducted under an inert atmosphere unless specified otherwise. All commercially purchased materials and solvents were used without further purification unless specified otherwise.

NMR spectra were recorded on a Varian VNMRs 500 (<sup>1</sup>H: 500 MHz, <sup>13</sup>C: 126 MHz) spectrometer and prepared in deuterated solvents such as CDCl<sub>3</sub> and DMSO-d<sub>6</sub>. <sup>1</sup>H and <sup>13</sup>C chemical shifts were recorded in parts per million (ppm). Multiplicity of <sup>1</sup>H-NMR peaks are indicated by s – singlet, d – doublet, dd – doublets of doublets, t – triplet, pt – pseudo triplet, q – quartet, m – multiplet and coupling constants are given in Hertz (Hz).

Electrospray ionisation – high resolution mass spectra (ESI-HRMS) were obtained using a Bruker Daltonics Apex III where Apollo ESI was used as the ESI source. Most analyses were conducted by Dr A. K. Abdul-Sada at Sussex. The molecular ion peaks [M]<sup>+</sup> were recorded as mass to charge (m/z) ratio. A portion of the high-resolution mass spectrometry (HRMS) measurements were also performed by the EPSRC National Mass Spectrometry Facility, University of Swansea.

LC-MS spectra were acquired using a Shimadzu LC-MS 2020, on a Gemini 5 μm C18 110 Å column and percentage purities were run over 30 minutes in water/acetonitrile with 0.1% formic acid (5 min at 5%, 5%-95% over 20 min, 5 min at 95%) with the UV detector at 254 nm. Purifications were performed by flash chromatography on silica gel columns or C18 columns using a Combi flash RF 75 PSI, ISCO unit.

## General Procedure

To a stirred solution of the appropriate 1,4-benzodiazepine or 1,3,4-benzotriazepine (0.030 mmol – 1.00 mmol, 1 eq.) and diaryliodonium salt (0.033 – 1.10 mmol, 1.1 eq.) in DCE (5 – 10 mL) was added 25% w/w NH<sub>3</sub> solution (aq. 5 – 10 mL) and the reaction mixture was stirred for 30 min. (unless stated otherwise). Upon completion, the reaction mixture was diluted with dichloromethane (3x15 mL) and the layers were separated. Combined organic layers were dried (MgSO<sub>4</sub>), concentrated under reduced pressure and purified by column chromatography, hexane/ethyl acetate (80:20 to 30:70).

### 1-(4-Nitrophenyl)-5-methyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one (**3a**)

The product was obtained as white solid (0.60 mmol scale, 170 mg, 96%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 8.27 – 8.21 (m, ArH, 2H), 7.62 (dd, <sup>3</sup>J<sub>HH</sub> = 7.5, 1.5 Hz, ArH, 1H), 7.40 – 7.35 (m, ArH, 3H), 7.34 – 7.29 (m, ArH, 1H), 6.82 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, ArH, 1H), 4.70 (d, <sup>2</sup>J<sub>HH</sub> = 10.5 Hz, COCH<sub>2</sub>, 1H), 3.83 (d, <sup>2</sup>J<sub>HH</sub> = 10.5 Hz, COCH<sub>2</sub>, 1H), 2.62 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 170.1 (C=O), 168.1 (C=N), 146.7 (ArC), 146.0 (ArC), 140.8 (ArC), 131.4 (ArC), 131.3 (ArC), 128.7 (ArC x 2), 127.8 (ArC), 125.9 (ArC), 125.1 (ArC), 124.5 (ArC x 2), 56.6 (COCH<sub>2</sub>), 25.5 (CH<sub>3</sub>). HRMS-ESI (m/z) calculated for C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub> [H]<sup>+</sup>: 296.1030, found: 296.1033. LCMS purity (UV) = 100%, t<sub>R</sub> 8.10 min.

### 1-(4-Nitrophenyl)-5-(propan-2-yl)-1,3-dihydro-2H-1,4-benzodiazepin-2-one (**3b**)

The product was obtained as a white solid (0.52 mmol scale, 166 mg, 99%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 8.27 – 8.20 (m, ArH, 2H), 7.59 (dd, <sup>3</sup>J<sub>HH</sub> = 7.5, 2.0 Hz, ArH, 1H), 7.39 – 7.35 (m, ArH, 3H), 7.34 – 7.30 (m, ArH, 1H), 6.83 (dd, <sup>3</sup>J<sub>HH</sub> = 8.0, 1.5 Hz, ArH, 1H), 4.72 (d, <sup>2</sup>J<sub>HH</sub> = 10.5 Hz, COCH<sub>2</sub>, 1H), 3.82 (d, <sup>2</sup>J<sub>HH</sub> = 10.5 Hz, COCH<sub>2</sub>, 1H), 3.34 – 3.25 (m, 1H), 1.35 (d, <sup>3</sup>J<sub>HH</sub> = 7.0 Hz, CNCHC<sub>2</sub>H<sub>5</sub>, 3H), 1.11 (d, <sup>3</sup>J<sub>HH</sub> = 7.0 Hz, CNCHC<sub>2</sub>H<sub>5</sub>, 3H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 176.9 (C=O), 168.7 (C=N), 146.7 (ArC), 145.9 (ArC), 141.5 (ArC), 131.6 (ArC), 130.9 (ArC), 128.3 (ArC x 2), 127.0 (ArC), 126.0 (ArC), 125.0 (ArC), 124.5 (ArC x 2), 56.5 (COCH<sub>2</sub>), 35.6 (CNCHC<sub>2</sub>H<sub>5</sub>), 22.0 (CNCHC<sub>2</sub>H<sub>5</sub>), 19.2 (CNCHC<sub>2</sub>H<sub>5</sub>). HRMS-ESI (m/z) calculated for C<sub>18</sub>H<sub>17</sub>N<sub>3</sub>O<sub>3</sub> [H]<sup>+</sup>: 324.1270, found: 324.1281. LCMS purity (UV) = 96 %, t<sub>R</sub> 18.73 min.

### 1-(4-Nitrophenyl)-3-(propan-2-yl)-5-(propan-2-yl)-1,3-dihydro-2H-1,4-benzodiazepin-2-one (**3c**)

The product was obtained as white solid (0.25 mmol scale, 91 mg, 99%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 8.25 – 8.18 (m, ArH, 2H), 7.61 (dd, <sup>3</sup>J<sub>HH</sub> = 8.0, 1.5 Hz, ArH, 1H), 7.39 – 7.24 (m, ArH, 4H), 6.85 (dd, <sup>3</sup>J<sub>HH</sub> = 8.0, 1.5 Hz, ArH, 1H), 3.27 (hept, <sup>3</sup>J<sub>HH</sub> = 7.0 Hz, CNCHC<sub>2</sub>H<sub>5</sub>, 1H), 3.12 (d, <sup>3</sup>J<sub>HH</sub> = 9.5 Hz, COCHC<sub>2</sub>H<sub>5</sub>, 1H), 2.72 – 2.61 (m, COCHC<sub>2</sub>H<sub>5</sub>, 1H), 1.33 (d, <sup>3</sup>J<sub>HH</sub> = 7.0 Hz, CNCHC<sub>2</sub>H<sub>5</sub>, 3H), 1.07 (d, <sup>3</sup>J<sub>HH</sub> = 7.0 Hz, CNCHC<sub>2</sub>H<sub>5</sub>, 3H), 1.05 – 1.02 (m, COCHC<sub>2</sub>H<sub>5</sub>, 6H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 173.9 (C=O), 168.3 (C=N), 147.4 (ArC), 145.7 (ArC), 141.1 (ArC), 131.9 (ArC), 130.6 (ArC), 128.4 (ArC x 2), 126.8 (ArC), 125.7 (ArC), 125.1

(ArC), 124.4 (ArC x 2), 69.3 (COCHCH<sub>2</sub>H<sub>6</sub>), 35.5 (CNCHCH<sub>3</sub>CH<sub>3</sub>), 22.2 (COCHCH<sub>2</sub>H<sub>6</sub>), 21.9 (CNCH<sub>2</sub>H<sub>6</sub>), 20.1, (CNCH<sub>2</sub>H<sub>6</sub>) 19.3 (COCHCH<sub>2</sub>H<sub>6</sub>), 18.7 (COCHCH<sub>2</sub>H<sub>6</sub>). HRMS-ESI (m/z) calculated for C<sub>21</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub> [+H]<sup>+</sup>: 366.1812, found: 366.1816. LCMS purity (UV) = 95%, tR 23.47 min.

**1-(4-Nitrophenyl)-5-phenyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one (3d)**

The product was obtained as white solid (0.60 mmol scale, 176 mg, 82%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 8.30 – 8.23 (m, ArH, 2H), 7.77 – 7.71 (m, ArH, 2H), 7.55 – 7.51 (m, ArH, 1H), 7.49 – 7.45 (m, ArH, 3H), 7.45 – 7.41 (m, ArH, 3H), 7.29 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, ArH, 1H), 6.94 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, ArH, 1H), 4.96 (d, <sup>2</sup>J<sub>HH</sub> = 10.5 Hz, COCH<sub>2</sub>, 1H), 4.03 (d, <sup>2</sup>J<sub>HH</sub> = 10.5 Hz, COCH<sub>2</sub>, 1H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 170.3 (C=O), 168.3 (C=N), 146.7 (ArC), 146.0 (ArC), 142.7 (ArC), 138.4 (ArC), 131.4 (ArC), 130.8 (ArC), 130.4 (ArC), 130.3 (ArC), 129.4 (ArC x 2), 128.5 (ArC x 2), 128.4 (ArC x 2), 125.4 (ArC), 125.0 (ArC), 124.5 (ArC x 2), 57.4 (COCH<sub>2</sub>). HRMS-ESI (m/z) calculated for C<sub>21</sub>H<sub>15</sub>N<sub>3</sub>O<sub>3</sub> [+H]<sup>+</sup>: 358.1186, found: 358.1187. LCMS purity (UV) = 95%, tR 18.35 min.

**1-(4-Nitrophenyl)-3-benzyl-5-phenyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one (3e)**

The product was obtained as white solid (0.40 mmol scale, 140 mg, 78%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 8.27 – 8.21 (m, ArH, 2H), 7.67 (d, <sup>3</sup>J<sub>HH</sub> = 7.5 Hz, ArH, 2H), 7.52 – 7.48 (m, 1H), 7.47 – 7.43 (m, ArH, 2H), 7.41 – 7.37 (m, ArH, 5H), 7.36 – 7.30 (m, ArH, 3H), 7.25 – 7.21 (m, ArH, 2H), 6.90 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, ArH, 1H), 4.01 (dd, J = 7.5, 6.0 Hz, COCHCH<sub>2</sub>, 1H), 3.68 (dd, <sup>2,3</sup>J<sub>HH</sub> = 14.0, 6.0 Hz, COCHCH<sub>2</sub>, 1H), 3.62 (dd, <sup>2,3</sup>J<sub>HH</sub> = 14.0, 7.5 Hz, COCHCH<sub>2</sub>, 1H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 168.8 (C=O), 168.5 (C=N), 147.1 (ArC), 145.9 (ArC), 142.1 (ArC), 138.9 (ArC), 138.4 (ArC), 131.5 (ArC), 130.8 (ArC), 130.5 (ArC), 130.3 (ArC), 130.0 (ArC x 2), 129.5 (ArC x 2), 128.6 (ArC x 2), 128.5 (ArC x 2), 128.3 (ArC x 2), 126.3 (ArC), 125.3 (ArC), 125.1 (ArC), 124.5 (ArC x 2), 65.6 (COCHCH<sub>2</sub>), 37.9 (COCHCH<sub>2</sub>). HRMS-ESI (m/z) calculated for C<sub>28</sub>H<sub>21</sub>N<sub>3</sub>O<sub>3</sub> [+H]<sup>+</sup>: 448.1656, found: 448.1669. LCMS purity (UV) = 99%, tR 20.81 min.

**7-Chloro-1-(4-nitrophenyl)-3-benzyl-5-phenyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one (3f)**

The product was obtained as white solid (0.15 mmol scale, 54 mg, 75%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 8.26 (d, <sup>3</sup>J<sub>HH</sub> = 8.5 Hz, ArH, 2H), 7.66 (d, <sup>3</sup>J<sub>HH</sub> = 7.5 Hz, ArH, 2H), 7.57 – 7.50 (m, ArH, 1H), 7.51 – 7.44 (m, ArH, 2H), 7.42 – 7.36 (m, ArH, 3H), 7.34 – 7.30 (m, ArH, 2H), 7.26 (s, ArH, 3H), 7.17 (d, J = 8.7 Hz, ArH, 1H), 6.85 (d, <sup>3</sup>J<sub>HH</sub> = 8.5 Hz, ArH, 1H), 3.99 (dd, J = 7.5, 6.0 Hz, COCHCH<sub>2</sub>, 1H), 3.70 – 3.57 (m, COCHCH<sub>2</sub>, 2H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 168.4 (C=O), 167.2 (C=N), 146.6 (ArC), 146.1 (ArC), 140.6 (ArC), 138.6 (ArC), 137.7 (ArC), 131.7 (ArC), 131.1 (ArC), 129.9 (ArC x 2), 129.8 (ArC), 129.5 (ArC x 2), 128.7 (ArC x 2), 128.6 (ArC x 2), 128.3 (ArC x 2), 126.5 (ArC), 126.4 (ArC), 126.2 (ArC), 124.6 (ArC x 2), 119.3 (ArC), 65.8 (COCHCH<sub>2</sub>), 37.9 (COCHCH<sub>2</sub>). HRMS-ESI (m/z) calculated for C<sub>28</sub>H<sub>20</sub>ClN<sub>3</sub>O<sub>3</sub> [+H]<sup>+</sup>: 482.1266, found: 482.1286. LCMS purity (UV) = 95%, tR 19.71 min.

**1-(4-Nitrophenyl)-3-benzyl-5-(pyridine-2-yl)-1,3-dihydro-2H-1,4-benzodiazepin-2-one (3g)**

The product was obtained as white solid (0.11 mmol scale, 38 mg, 77%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 8.67 – 8.62 (m, ArH, 1H), 8.15 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, ArH, 2H), 8.18 – 8.12 (m, ArH, 1H), 7.88 – 7.81 (m, ArH, 1H), 7.44 – 7.42 (m, ArH, 2H), 7.42 – 7.37 (m, ArH, 4H), 7.35 – 7.26 (m, ArH, 3H), 7.25 – 7.21 (m, ArH, 2H), 6.89 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, ArH, 1H), 4.10 (dd, <sup>3</sup>J<sub>HH</sub> = 8.0, 6.0 Hz, COCHCH<sub>2</sub>, 1H), 3.70 (dd, <sup>2,3</sup>J<sub>HH</sub> = 14.0, 7.0 Hz, COCHCH<sub>2</sub>, 1H), 3.62 (dd, <sup>2,3</sup>J<sub>HH</sub> = 14.0, 7.5 Hz, COCHCH<sub>2</sub>, 1H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 168.6 (C=O), 167.6 (C=N), 155.9 (ArC), 148.7 (ArC), 147.1 (ArC), 145.9 (ArC), 141.9 (ArC), 138.9 (ArC), 136.8 (ArC), 131.4 (ArC), 130.8 (ArC), 129.9 (ArC x 2), 128.8 (ArC x 2), 128.3 (ArC x 2), 126.3 (ArC), 125.2 (ArC), 125.1 (ArC), 124.8

(ArC x 2), 124.4 (ArC x 2), 123.8 (ArC), 65.8 (COCHCH<sub>2</sub>), 37.8 (COCHCH<sub>2</sub>). HRMS-ESI (m/z) calculated for C<sub>27</sub>H<sub>20</sub>N<sub>4</sub>O<sub>3</sub> [+H]<sup>+</sup>: 449.1608, found: 449.1617. LCMS purity (UV) = 99%, tR 20.81 min.

**1-(4-Nitrophenyl)-3-benzyl-5-(2'-fluorobiphenyl-2-yl)-1,3-dihydro-2H-1,4-benzodiazepin-2-one (3h)**

The product was obtained as white solid (0.03 mmol scale, 11 mg, 70%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 8.14 (d, <sup>3</sup>J<sub>HH</sub> = 8.5 Hz, ArH, 2H), 7.57 – 7.52 (m, ArH, 1H), 7.51 – 7.45 (m, ArH, 1H), 7.42 (d, <sup>3</sup>J<sub>HH</sub> = 7.5 Hz, ArH, 1H), 7.38 (d, <sup>3</sup>J<sub>HH</sub> = 7.5 Hz, ArH, 1H), 7.32 – 7.27 (m, ArH, 7H), 7.26 – 7.19 (m, ArH, 2H), 7.15 – 7.09 (m, ArH, 1H), 7.06 (d, <sup>3</sup>J<sub>HH</sub> = 7.5 Hz, ArH, 1H), 7.00 – 6.93 (m, ArH, 3H), 6.65 (d, <sup>3</sup>J<sub>HH</sub> = 8.5 Hz, ArH, 1H), 3.80 (dd, <sup>3</sup>J<sub>HH</sub> = 8.0, 5.5 Hz, COCHCH<sub>2</sub>, 1H), 3.69 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, COCHCH<sub>2</sub>, 1H), 3.66 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, COCHCH<sub>2</sub>, 1H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 169.5 (C=O), 167.9 (C=N), 159.2 (d, <sup>1</sup>J<sub>FC</sub> = 247.5 Hz, ArC), 147.1 (ArC), 145.7 (ArC), 141.5 (ArC), 138.8 (ArC), 138.7 (ArC), 135.7 (ArC), 132.0 (d, <sup>3</sup>J<sub>FC</sub> = 3.5 Hz, ArC), 131.6 (ArC), 131.4 (ArC), 130.8 (ArC), 130.3 (ArC), 129.9 (ArC x 2), 129.8 (ArC), 129.5 (ArC), 129.2 (ArC), 128.9 (d, <sup>3</sup>J<sub>FC</sub> = 8.0 Hz, ArC), 128.5 (ArC x 2), 128.3 (ArC x 2), 128.1 (ArC), 126.3 (ArC), 125.2 (ArC), 124.8 (ArC), 124.5 (d, <sup>4</sup>J<sub>FC</sub> = 3.5 Hz, ArC), 124.2 (ArC x 2), 115.4 (d, <sup>2</sup>J<sub>FC</sub> = 22.0 Hz, ArC), 66.0 (COCHCH<sub>2</sub>), 37.8 (COCHCH<sub>2</sub>). HRMS-ESI (m/z) calculated for C<sub>34</sub>H<sub>24</sub>N<sub>3</sub>O<sub>3</sub> [+H]<sup>+</sup>: 542.1874, found: 542.1881. LCMS purity (UV) = 93%, tR 23.27 min.

**1-(2,4,6-Trimethylphenyl)-5-phenyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one (3i)**

The reaction was run for 8 hours. The product was obtained as white solid (1.00 mmol scale, 181 mg, 51%). <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 7.60 – 7.57 (m, ArH, 2H), 7.54 – 7.46 (m, ArH, 4H), 7.32 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, ArH, 1H), 7.27 – 7.23 (m, ArH, 1H), 7.10 – 7.07 (m, ArH, 1H), 6.88 (s, ArH, 1H), 6.78 (d, <sup>3</sup>J<sub>HH</sub> = 8.0, 1.1 Hz, ArH, 1H), 4.70 (d, <sup>2</sup>J<sub>HH</sub> = 10.0 Hz, COCH<sub>2</sub>, 1H), 4.04 (d, <sup>2</sup>J<sub>HH</sub> = 10.0 Hz, COCH<sub>2</sub>, 1H), 2.26 (s, CH<sub>3</sub>, 3H), 2.24 (s, CH<sub>3</sub>, 3H), 1.61 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C-NMR (126 MHz) DMSO-*d*<sub>6</sub>: δ = 170.3 (C=O), 167.5 (C=N), 142.2 (ArC), 138.9 (ArC), 137.9 (ArC), 137.0 (ArC), 136.2 (ArC), 134.7 (ArC), 132.3 (ArC), 130.9 (ArC), 130.0 (ArC x 2), 129.6 (ArC x 2), 129.5 (ArC), 128.9 (ArC x 2), 128.7 (ArC), 124.4 (ArC), 122.1 (ArC), 57.3 (COCH<sub>2</sub>), 21.0 (CH<sub>3</sub>), 18.5 (CH<sub>3</sub>), 17.5 (CH<sub>3</sub>). HRMS-ESI (m/z) calculated for C<sub>24</sub>H<sub>22</sub>N<sub>2</sub>O [+H]<sup>+</sup>: 355.1805, found: 355.1804. LCMS purity (UV) = 97%, tR 21.13 min.

**1-(2-Bromophenyl)-5-phenyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one (3j)**

The reaction was run for 8 hours. The product was obtained as white solid (1.00 mmol scale, 31 mg, 8%). <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 7.84 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, ArH, 1H), 7.70 (d, <sup>3</sup>J<sub>HH</sub> = 8.0 Hz, ArH, 1H), 7.66 – 7.59 (m, ArH, 4H), 7.52 – 7.46 (m, ArH, 3H), 7.41 – 7.37 (m, ArH, 1H), 7.32 (dd, J = 7.8, 1.7 Hz, ArH, 1H), 7.27 (d, <sup>3</sup>J<sub>HH</sub> = 7.0 Hz, ArH, 1H), 6.92 – 6.83 (m, ArH, 1H), 4.69 (d, <sup>2</sup>J<sub>HH</sub> = 10.5 Hz, COCH<sub>2</sub>, 1H), 4.01 (d, <sup>2</sup>J<sub>HH</sub> = 10.5 Hz, COCH<sub>2</sub>, 1H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 170.7 (C=O), 168.9 (C=N), 142.0 (ArC), 138.9 (ArC), 138.6 (ArC), 134.3 (ArC), 133.7 (ArC), 132.0 (ArC), 131.0 (ArC), 130.9 (ArC), 130.8 (ArC), 129.9 (ArC x 2), 129.8 (ArC), 129.1 (ArC), 128.8 (ArC x 2), 124.7 (ArC), 123.0 (ArC), 121.5 (ArC), 57.0 (COCH<sub>2</sub>). HRMS-ESI (m/z) calculated for C<sub>21</sub>H<sub>15</sub>BrN<sub>2</sub>O [+H]<sup>+</sup>: 391.0441, found: 391.0457. LCMS purity (UV) = 93%, tR 15.23 min.

**1-(3'-Trifluoromethylphenyl)-5-phenyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one (3k)**

The reaction was run for 8 hours. The product was obtained as white solid (0.50 mmol scale, 80 mg, 42%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 7.72 (d, <sup>3</sup>J<sub>HH</sub> = 7.5 Hz, ArH, 2H), 7.60 – 7.55 (m, ArH, 2H), 7.53 (d, <sup>3</sup>J<sub>HH</sub> = 8.5 Hz, ArH, 2H), 7.49 – 7.45 (m, ArH, 2H), 7.44 – 7.38 (m, ArH, 3H), 7.25 – 7.20 (m, ArH, 1H), 6.92 (d, <sup>3</sup>J<sub>HH</sub> = 8.5 Hz, ArH, 1H), 4.95 (d, <sup>2</sup>J<sub>HH</sub> = 10.5 Hz, 1H), 4.02 (d, <sup>2</sup>J<sub>HH</sub> = 10.5 Hz, 1H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 170.3 (C=O), 168.3 (C=N), 143.1 (ArC), 141.3 (ArC), 138.6 (ArC), 132.0 (q, <sup>2</sup>J<sub>FC</sub> = 29.9

Hz, ArC), 131.9 (ArC), 131.6 (ArC), 131.4 (ArC), 130.7 (ArC), 130.4 (ArC), 129.8 (ArC x 2), 129.5 (ArC), 128.5 (ArC x 2), 125.2 (q,  $^3J_{\text{FC}} = 3.5$  Hz, ArC), 124.8 (ArC), 124.8 (ArC), 123.5 (q,  $^1J_{\text{FC}} = 273.0$  Hz, ArC), 124.2 (q,  $^3J_{\text{FC}} = 3.5$  Hz, ArC), 57.3 (COCH<sub>2</sub>). HRMS-ESI (m/z) calculated for C<sub>22</sub>H<sub>15</sub>F<sub>3</sub>N<sub>2</sub>O [H]<sup>+</sup>: 381.1209, found: 381.1208. LCMS purity (UV) = 96%, t<sub>R</sub> 21.35 min.

### 1-Phenyl-5-phenyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one (3I)

The reaction was run for 8 hours. The product was obtained as white solid (0.50 mmol scale, 14 mg, 9%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 7.72 (d,  $^3J_{\text{HH}} = 7.5$  Hz, ArH, 2H), 7.54 – 7.49 (m, ArH, 1H), 7.47 (d,  $^3J_{\text{HH}} = 7.5$  Hz, ArH, 2H), 7.43 – 7.38 (m, ArH, 2H), 7.37 – 7.30 (m, ArH, 2H), 7.24 – 7.21 (m, ArH, 3H), 7.20 – 7.16 (m, ArH, 1H), 6.97 (d,  $^3J_{\text{HH}} = 8.5$  Hz, ArH, 1H), 4.96 (d,  $^2J_{\text{HH}} = 10.5$  Hz, ArH, COCH<sub>2</sub>, 1H), 4.01 (d,  $^2J_{\text{HH}} = 10.5$  Hz, COCH<sub>2</sub>, 1H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 170.7 (C=O), 168.3 (C=N), 146.5 (ArC), 143.3 (ArC), 140.7 (ArC), 138.6 (ArC), 131.3 (ArC), 130.7 (ArC), 130.3 (ArC), 129.6 (ArC x 2), 129.3 (ArC x 2), 128.4 (ArC x 2), 128.3 (ArC x 2), 127.5 (ArC), 124.7 (ArC), 124.2 (ArC), 57.2 (COCH<sub>2</sub>). HRMS-ESI (m/z) calculated for C<sub>21</sub>H<sub>16</sub>N<sub>2</sub>O [H]<sup>+</sup>: 313.1335, found: 313.1338. LCMS purity (UV) = 90%, t<sub>R</sub> 16.10 min.

### 1-(4-Nitrophenyl)-3-(4-nitrophenyl)-5-phenyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one (7)

The product was obtained as white solid (0.60 mmol scale, 2 eq. of diaryliodonium triflate, 146 mg, 51%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 8.34 – 8.22 (m, ArH, 4H), 7.85 – 7.79 (m, ArH, 2H), 7.77 – 7.71 (m, ArH, 2H), 7.74 – 7.57 (m, ArH, 3H), 7.59 – 7.50 (m, ArH, 3H), 7.41 – 7.32 (m, ArH, 2H), 7.07 – 7.02 (m, ArH, 1H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 166.0 (C=O), 158.3 (C=N), 149.1 (ArC), 146.7 (ArC), 145.6 (ArC), 144.1 (ArC), 143.3 (ArC), 135.0 (ArC), 132.6 (ArC), 131.5 (ArC), 129.9 (ArC), 129.6 (ArC), 129.5 (ArC x 2), 128.9 (ArC x 2), 126.8 (ArC x 2), 126.3 (ArC), 125.4 (ArC), 124.5 (ArC x 2), 124.3 (ArC x 2), 121.3 (ArC x 2). HRMS-ESI (m/z) calculated for C<sub>26</sub>H<sub>17</sub>N<sub>5</sub>O<sub>5</sub> [H]<sup>+</sup>: 480.1230, found: 480.1245. LCMS purity (UV) = 95%, t<sub>R</sub> 18.35 min.

### 1,1'-Oxybis(4-nitrobenzene) (10)

To a solution of (4-nitrophenyl)phenyliodonium triflate (30 mg, 0.06 mmol) in DCE (1 mL) was added sodium hydroxide (aq., 1 mL) and stirred for 20 min. at room temperature. Upon completion, the reaction was diluted with dichloromethane (5 mL x 3) and the layers were separated. Combined organic layers were dried (MgSO<sub>4</sub>) and concentrated under reduced pressure to afford the product as a white powder (7 mg, 43%). <sup>1</sup>H-NMR (500 MHz) CDCl<sub>3</sub>: δ = 8.33 – 8.27 (m, ArH, 4H), 7.19 – 7.14 (m, ArH, 4H). <sup>13</sup>C-NMR (126 MHz) CDCl<sub>3</sub>: δ = 160.6 (ArC x 2), 144.2 (ArC x 2), 126.2 (ArC x 4), 119.3 (ArC x 4). HRMS-ESI (m/z) calculated for C<sub>12</sub>H<sub>8</sub>N<sub>2</sub>O<sub>5</sub> [H]<sup>+</sup>: 261.0511, found: 261.0513.

### Funding Information

R.K. is funded by an EPSRC/AZ funded PhD studentship (EP/M507568/1) with additional support from AstraZeneca [14550001 (SME)] and Tocris Biosciences.

### Acknowledgment

We thank Dr. Alaa Abdul-Sada (Sussex) and the EPSRC UK National Mass Spectrometry Facility at Swansea University for HR-MS measurements.

### Supporting Information

YES (this text will be updated with links prior to publication)

### Primary Data

NO (this text will be deleted prior to publication)

### References and Notes

- Filippakopoulos, P.; Qi, J.; Picaud, S.; Shen, Y.; Smith, W. B.; Fedorov, O.; Morse, E. M.; Keates, T.; Hickman, T. T.; Felletar, I.; Philpott, M.; Munro, S.; McKeown, M. R.; Wang, Y.; Christie, A. L.; West, N.; Cameron, M. J.; Schwartz, B.; Heightman, T. D.; La Thangue, N.; French, C. A.; Wiest, O.; Kung, A. L.; Knapp, S.; Bradner, J. E. *Nature* **2010**, 468, 1067.
- Field, G. F.; Zally, W. J.; Sternbach, L. H. *J. Am. Chem. Soc.* **1967**, 80, 332.
- Ellmann, J. A. *Acc. Chem. Res.* **1996**, 29, 132.
- Nadin, A.; Sa'nchez Lo'pez, J. M.; Owens, A. P.; Howells, D. M.; Talbot, A. C.; Harrison, T. *J. Org. Chem.* **2003**, 68, 2844.
- Baud, M. G.; Lin-Shiao, E.; Cardote, T.; Tallant, C.; Pschibul, A.; Chan, K. H.; Zengerle, M.; Garcia, J. R.; Kwan, T. T.; Ferguson, F. M.; Ciulli, A. *Science* **2014**, 346, 638.
- Baud, M. G.; Lin-Shiao, E.; Zengerle, M.; Tallant, C.; Ciulli, A. *J. Med. Chem.* **2016**, 59, 1492.
- Mirguet, O.; Gosmini, R.; Toum, J.; Clement, C. A.; Barnathan, M.; Brusq, J. M.; Mordaunt, J. E.; Grimes, R. M.; Crowe, M.; Pineau, O.; Ajakane, M.; Daugan, A.; Jeffrey, P.; Cutler, L.; Haynes, A. C.; Smithers, N. N.; Chung, C. W.; Bamborough, P.; Uings, I. J.; Lewis, A.; Witherington, J.; Parr, N.; Prinjha, R. K.; Nicodeme, E. *J. Med. Chem.* **2013**, 56, 7501.
- Liu, J. J.; Higgins, B.; Ju, G.; Kolinsky, K.; Luk, K. C.; Packman, K.; Pizzolato, G.; Ren, Y.; Thakkar, K.; Tovar, C.; Zhang, Z.; Wovkulich, P. M. *ACS Med. Chem. Lett.* **2013**, 4, 259.
- Evans, B.; Rittle, K.; Bock, M.; DiPardo, R.; Freidinger, R.; Whitter, W.; Lundell, G.; Veber, D.; Anderson, P.; Chang, R.; Lotti, V.; Cerino, D.; Chen, T.; Kling, P.; Kunkel, K.; Springer, J.; Hirshfield, J. *J. Med. Chem.* **1988**, 31, 2235.
- Filippakopoulos, P.; Picaud, S.; Fedorov, O.; Keller, M.; Wrobel, M.; Morgenstern, O.; Bracher, F.; Knapp, S. *Bioorg. Med. Chem.* **2012**, 20, 1878.
- Smith, S. G.; Sanchez, R.; Zhou, M.-M. *Chemistry & Biology* **2014**, 573.
- Filippakopoulos, P.; Knapp, S. *Nat Rev Drug Discov* **2014**, 13, 337.
- Carter, M. C.; Alber, D. G.; Baxter, R. C.; Bithell, S. K.; Budworth, J.; Chubb, A.; Cockerill, G. S.; Dowdell, V. C. L.; Henderson, E. A.; Keegan, S. J.; Kelsey, R. D.; Lockyer, M. J.; Stables, J. N.; Wilson, L. J.; Powell, K. L. *J. Med. Chem.* **2006**, 49, 2311.
- Ghelani, S. M.; Naliapara, Y. T. *J. Heterocyclic Chem.* **2016**, 53, 1795.
- Abdelkafi, H.; Cintrat, J. C. *Sci. Rep.* **2015**, 5, 12131.
- Kaur, N. *Int. J. Pharm. Bio. Sci.* **2013**, 485.
- Hu, X.; Dong, Y.; Liu, G. *Mol. Divers.* **2015**, 19, 695.
- Spencer, J.; Rathnam, R.; Chowdhry, B. *Fut. Med. Chem.* **2010**, 2, 1441.
- Dzladulewicz, E. K.; Brown, M. C.; Dunstan, A. R.; Lee, W.; Said, N. B.; Garratt, P. J. *Bioorg. Med. Chem. Lett.* **1999**, 9, 463.
- McDonald, I.; Austin, C.; Buck, I.; Dunstone, D.; Gaffen, J.; Griffin, E.; Harper, E.; Hull, R.; Kalindjian, S.; Linney, I.; Low, C.; Patel, D.; Pether, M.; Raynor, M.; Roberts, S.; Shaxted, M.; Spencer, J.; Steel, K.; Sykes, D.; Wright, P.; Xun, W. *J. Med. Chem.* **2007**, 50, 4789.
- Cuny, G.; Bois-Choussy, M. I.; Zhu, J. *J. Am. Chem. Soc.* **2004**, 126, 14475.
- Klapars, A.; Antilla, J. C.; Huang, X.; Buchwald, S. L. *J. Am. Chem. Soc.* **2001**, 123, 7727.

- (23) Talukdar, D.; Das, G.; Thakur, S.; Karak, N.; Thakur, A. J. *Catal. Commun.* **2015**, *59*, 238.
- (24) Strittmatter, S. M. *Nat. Med.* **2014**, *20*, 590.
- (25) Corsello, S. M.; Bittker, J. A.; Liu, Z.; Gould, J.; McCarren, P.; Hirschman, J. E.; Johnston, S. E.; Vrcic, A.; Wong, B.; Khan, M.; Asiedu, J.; Narayan, R.; Mader, C. C.; Subramanian, A.; Golub, T. R. *Nat. Med.* **2017**, *23*, 405.
- (26) Spencer, J.; Rathnam, R. P.; Harvey, A. L.; Clements, C. J.; Clark, R. L.; Barrett, M. P.; Wong, P. E.; Male, L.; Coles, S. J.; Mackay, S. P. *Bioorg. Med. Chem.* **2011**, *19*, 1802.
- (27) Clark, R. L.; Clements, C. J.; Barrett, M. P.; Mackay, S. P.; Rathnam, R. P.; Owusu-Dapaah, G.; Spencer, J.; Huggan, J. K. *Bioorg. Med. Chem.* **2012**, *20*, 6019.
- (28) Spencer, J.; Chowdhry, B. Z.; Mallet, A. I.; Rathnam, R. P.; Adatia, T.; Bashall, A.; Rominger, F. *Tetrahedron* **2008**, *64*, 6082.
- (29) Khan, R.; Felix, R.; Kemmitt, P. D.; Coles, S. J.; Day, I. J.; Tizzard, G. J.; Spencer, J. *Adv. Synth. Catal.* **2016**, *358*, 98.
- (30) Merritt, E. A.; Olofsson, B. *Angew. Chem. Int. Ed. Engl.* **2009**, *48*, 9052.
- (31) Bielawski, M.; Aili, D.; Olofsson, B. *J. Org. Chem.* **2008**, *73*, 4602.
- (32) Yusubov, M. S.; Maskae, A. V.; Zhdankin, V. V. *Arkivoc* **2011**, *1*, 370.
- (33) Ghosh, R.; Olofsson, B. *Org. Lett.* **2014**, *16*, 1830.
- (34) Tinnis, F.; Stridfeldt, E.; Lundberg, H.; Adolfsson, H.; Olofsson, B. *Org. Lett.* **2015**, *17*, 2688.
- (35) Malmgren, J.; Santoro, S.; Jalalian, N.; Himo, F.; Olofsson, B. *Chem. Eur. J.* **2013**, *19*, 10334.
- (36) Gonda, Z.; Novak, Z. *Chem. Eur. J.* **2015**, *21*, 16801.
- (37) Lanning, M.; Fletcher, S. *Fut. Med. Chem.* **2013**, *5*, 2157.
- (38) Azzarito, V.; Long, K.; Murphy, N. S.; Wilson, A. J. *Nat. Chem.* **2013**, *5*, 161.
- (39) Jalalian, N.; Ishikawa, E. E.; Silva Jr., L. F.; Olofsson, B. *Org. Lett.* **2011**, *13*, 1552.
- (40) Coles, S. J.; Gale, P. A. *Chem. Sci.*, **2012**, *3*, 683.